

S.P.Novikov, M.A.Shubin

# The Morse Theory and von Neuman Topological Invariants for Non-Simply-Connected Manifolds<sup>1</sup>

The real numbers  $\bar{b}_p(X)$  were invented first by Atiyah (see[1]) for compact non-simply-connected  $n$ -dimensional manifolds  $X$  through the  $L_2$ -cohomology of its universal covering  $M$ . More general numbers  $\bar{b}_p^\chi$  were defined by Singer (see [2]) for every representation  $\chi : \pi_1(X) \rightarrow A$  where  $A$  is a von Neuman algebra with finite normalized trace (the Atiyah numbers  $\bar{b}_p(X)$  correspond to the regular representation  $\chi$  of the group  $\pi_1(X) = \Gamma$  into the space  $l_2(\Gamma)$ . It was found (see [3]) that in the classical Morse inequalities on the manifold  $X$  one can replace ordinary Betti numbers  $b_p$  by the numbers  $\bar{b}_p^\chi$  for every  $\chi$ . In some cases von Neuman-Morse Inequalities are stronger than the ordinary Morse Inequalities. Sometimes they allow to establish non-triviality of the  $L_2$ -cohomology for coverings of non-simply-connected manifolds.

It was established in the works [4, 5] that for  $M = H^n$  (Hyperbolic or Lobachevski Space) and  $M$ -strictly Convex Domain with Bergman metric we have  $\bar{b}_p = 0$  for  $p \neq n/2$ . In the cases where  $\bar{b}_p = 0$  and the Spectrum of all Laplacians on the spaces of  $p$ -forms are separated from zero and or spectrum touches zero, we define a von Neuman Analog of the Ray-Singer Torsion  $\bar{R}$ .

The fact that the Spectrum of  $\Delta_p$  touches zero, does not depend on Riemannian Metric on  $X$ . For  $p = 0$  such event takes place (see [6]) if and only if  $\Gamma$  is amenable. For  $M = H^{2k+1}$  the spectrum of  $\Delta_p$  touches zero for  $p = k, k + 1$  only (see [4]). The power of  $t$  of the power-like decay for  $t \rightarrow \infty$  of the  $\theta(t) = Tr_\Gamma \exp(-t\Delta_p)$  also does not depend on metric, or the power entering the power-like asymptotics for  $\lambda \rightarrow +0$  for the density of states  $N_p(\lambda) = Tr_\Gamma E_\lambda(\Delta_p)$ . Here  $E_\lambda(\Delta_p)$  is the spectral projector of the operator  $\Delta_p$ , and the trace  $Tr_\Gamma$  is defined by the integration along the fundamental

---

<sup>1</sup>translated from Russian, Uspekhi Math Nauk= Russian Math Surveys, Section "Mathematical Life in the USSR", Meetings of I.G.Petrovski Seminar, March 5, 1986 (the first meeting), 1986, vol 41, n 4, pp 222-223

domain of the Kernel, according to [1]. Using Variational Principle analogous to [7], we prove following

**Theorem 1** *Let  $N_p, \theta_p$  and  $N', \theta'$  correspond to two different Riemannian Metrics on the compact manifold  $X$  covered by  $M$ . There exists positive nonzero constant  $C$  such that*

$$N_p(C^{-1}\lambda) \leq N'_p(\lambda) \leq N(C\lambda), C^{-1}\theta_p(Ct) \leq \theta'(t) \leq C^{-1}(C^{-1}t)$$

*If  $\theta \leq o(t^{-\epsilon})$  for all  $p$  for  $t \rightarrow \infty$  where  $\epsilon > 0$  for all  $p = 0, \dots, n$ , the quantity  $\bar{R}$  is well-defined.*

**Conjecture:** The estimates  $\theta_p(t) = o(t^{-\epsilon_p}), \epsilon_p > 0$ , are valid allways if  $\bar{b}_p = 0$ .

For  $M = H^3$  we have  $\theta_1(t) \sim ct^{-1/2}$  –see [8]<sup>2</sup>. It allows to define  $\bar{R}$  for 3-manifolds of the constant negative curvature. However, as Senya Vishik communicated to us privately, in this case  $\bar{R} = 0$ . Probably, nontrivial von Neumann Torsion might appear for the nontrivial representations different from the regular one<sup>3</sup>.

## References

- [1] Atiyah M. Elliptic Operators, discrete groups and von Neuman Algebras, Asterisque, 1976, Vol 32-33, pp 43-72
- [2] Singer I. Some remarks on Operator Theory and Index Theory, Lecture Notes in Math., Vol 575, pp 128-137.

---

<sup>2</sup>Originally this result was extracted by the present author (S.Novikov), who worked some period studying Cosmological Models of the General Relativity in early 1970s, from the old famous work of physicist E.Lifshitz on time evolution of the completely isotropic Friedman Cosmological Model with 3-space sections of constant negative curvature (1946). Misha Shubin replaced it by the quotation to the more recent rigorous mathematical work [8] of Senya Vishik

<sup>3</sup>It turned out that Senya Vishik made mistake in his calculation: John Lott recalculated it later by the suggestion of the present author (S.Novikov) and found out that density of torsion is nonzero indeed  $\bar{R} \neq 0$  in this case, so it is proportional to the volume of fundamental domain with nonzero coefficient. M.Gromov and M.Farber proved later that these „Novikov-Shubin” power invariants are homotopy invariant. They pointed out that they appear as natural homological quantities for infinite-dimensional complexes.

- [3] Novikov S., Shubin M. Morse Inequalities and von Neuman Algebras, Uspekhi Math Nauk=Russian Math Surveys (in Russian), 1986, vol 41, n 4 (this work appeared soon in Russian and in English in the Journal Doklady Akademii Nauk=Soviet Math Doklady, 1986, vol 289 n 2 pp 289-292 )
- [4] Donnelly H. The differential form spectrum of hyperbolic space. Manuscripta Math., 1980-81, vol 33, nn 3/4, pp365-385
- [5] Donnelly H., Fefferman Ch.  $L_2$ -cohomology and index theorem for the Bergman metric. Ann Math. 1983, vol 118, n 3 pp 593-618
- [6] Brooks B. The fundamental Group and the spectrum of the Laplacian. Comm Math Helv. 1981, vol 56, 581-598
- [7] Bogorodskaya T., Shubin M. Variational Principle for the density of States of the Random Pseudodifferential Operators, Funktsional Anal Appl., 1983, vol 17, n 2, pp 66-67
- [8] Vishik S. Some Analogs of the Selberg  $\zeta$ -function. Funktsional Anal Appl., 1975, vol 9, n 3, pp 85-86